

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

Supplied in a tight, light-resistant container with a child-resistant cap.

THIS BOTTLE NOT TO BE DISPENSED TO THE CONSUMER.

Mfg. for: Cypress Pharmaceutical, Inc.
Madison, MS 39110

L591 Rev. 03/09



NDC 60258-415-16

CERON-DM

Syrup

Antihistamine/Decongestant
Antitussive

Each teaspoonful (5 mL) for oral
administration contains:

Chlorpheniramine Maleate . . . 4.0 mg
Phenylephrine HCl 12.5 mg
Dextromethorphan HBr 15.0 mg

Grape Flavor

Sugar Free/Alcohol Free

CYPRESS
PHARMACEUTICAL, INC.

16 fl oz (473 mL)

Drug Facts

Lift Here 

Active ingredients (in each 5 mL teaspoonful)	Purpose
Chlorpheniramine Maleate 4 mg.....	Antihistamine
Phenylephrine Hydrochloride 12.5 mg.....	Decongestant
Dextromethorphan Hydrobromide 15 mg	Antitussive

Uses temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes
- nasal congestion ■ reduces swelling of nasal passages ■ cough due to minor throat and bronchial irritation

Warnings

Do not exceed recommended dosage.

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis ■ glaucoma
- trouble urinating due to an enlarged prostate gland ■ heart disease
- high blood pressure
- thyroid disease ■ diabetes mellitus

Drug Facts (continued)

- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- excitability may occur, especially in children
- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of accidental overdose seek medical help or contact a Poison Control Center immediately.

Directions

Do not exceed recommended dosage.

Adults and children 12 years of age and older:	1 teaspoonful (5 mL) every 6 hours, not to exceed 4 teaspoonfuls in 24 hours.
--	---

Drug Facts (continued)

Children 6 to under 12 years of age:	1/2 teaspoonful every 6 hours, not to exceed 2 teaspoonfuls in 24 hours.
--------------------------------------	--

Children under 6 years of age:	Consult a physician.
--------------------------------	----------------------

Other information

Store at 68° - 77°F (20° - 25°C)

Inactive ingredients

FD&C Blue #1, FD&C Red #40, Glycerin, Grape Flavor, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

Questions? Comments?

1-800-856-4393

M-F: 8:00 a.m. - 5:00 p.m. CT

Serious side effects associated with use of this product may be reported to this number.

Manufactured for:
Cypress Pharmaceutical, Inc.
Madison, MS 39110

I297

Rev. 03/09